

L25 ANSWER 2 OF 4  
 ACCESSION NUMBER:  
 DOCUMENT NUMBER:  
 TITLE:

MEDLINE  
 97031814 MEDLINE  
 97031814 PubMed ID: 8877720 .  
 Redirecting circulating antibodies via **ligand-hapten conjugates** eliminates target cells  
 in vivo.

AUTHOR:

Lussow A R; Buelow R; Fanget L; Peretto S; Gao L; Pouletty P

CORPORATE SOURCE:  
 SOURCE:  
 IMMUNOLOGY,

SangStat Medical Corporation, Menlo Park, CA 94025, USA.  
 JOURNAL OF IMMUNOTHERAPY WITH EMPHASIS ON TUMOR

(1996 Jul) 19 (4) 257-65.  
 Journal code: 9418950. ISSN: 1067-5582.

PUB. COUNTRY:  
 DOCUMENT TYPE:  
 LANGUAGE:  
 FILE SEGMENT:  
 ENTRY MONTH:  
 ENTRY DATE:

United States  
 Journal; Article; (JOURNAL ARTICLE)  
 English  
 Priority Journals  
 199702  
 Entered STN: 19970219  
 Last Updated on STN: 19970219  
 Entered Medline: 19970204

AB The elimination of cell populations in vivo often relies on reagents that are self-limiting, are difficult to design and produce or contain highly toxic components. Here we describe a novel immunotherapy using molecules that combine a cell-specific ligand and a hapten binding to preexisting antibodies in serum. The F(ab')<sub>2</sub> fragment of a polyclonal anti-thymocyte globulin (ATG) preparation was used as a T-cell-specific ligand, and fluorescein isothiocyanate (FITC), as the hapten. Clearance of **ligand-hapten conjugates** from the circulation through formation of immune complexes was prevented through controlled synthesis of conjugates so that they contained one F(ab')<sub>2</sub> fragment and one FITC molecule. Administration of a single dose of F(ab')<sub>2</sub> or F(ab')<sub>2</sub>ATG-FITC into naive mice had no effect on the number of circulating T cells. In contrast, injection of F(ab')<sub>2</sub>ATG-FITC into mice with circulating anti-FITC antibodies resulted in the elimination of peripheral T cells. The reduction in cell numbers was equivalent to that obtained with a corresponding dose of intact ATG. Experiments in thymectomized mice demonstrated that the reduction of circulating T cells was due to target-cell elimination and not to immunomodulation or cellular sequestration. The adaptability of the model to other sources of effector antibodies and more useful ligands is discussed.